
LAND SUITABILITY MAP

CACAO

LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS

PROVINCE OF APAYAO






MAP INDEX

LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS

PROVINCE OF APAYAO



LAND SUITABILITY MAP FOR CACAO

LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS

APAYAO, CAR

EXTENT OF SUITABILITY FOR CACAO PRODUCTION BY MUNICIPALITY

						EX	PANSION	AREA (H	la)			-	CONF	LICT RES	OLUTION	(Ha)			TOTAL
MUNICIPALITY	EXISTI	NG CACA	.O (Ha)	TOTAL EXISTING AREA (Ha)	Coco	nut	Shrub unman		Grass unmar		Co	rn	_	oaddy, rigated	Man	ıgo	Other	crops	POTENTIAL EXPANSION
	S1	S2	S 3		S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	AREA (Ha)
CALANASAN	4	1	833	838	-	-	366	-	172	4	147	14	108	7	-	-	-	-	818
CONNER	76	-	66	142	286	15	1,720	332	4,234	1,601	2,362	124	612	87	10	-	-	-	11,384
FLORA	-	-	-	-	-	-	7,572	113	3,933	25	1,170	5	1,467	62	1	-	-	-	14,349
KABUGAO	-	-	21	21	-	-	3,072	66	287	38	884	-	137	-	-	-	-	-	4,485
LUNA	-	36	14	50	-	-	1,777	278	1,398	87	1,470	108	1,520	258	-	-	-	-	6,896
PUDTOL	48	-	25	73	-	-	3,541	102	1,559	15	1,163	393	994	564	-	-	-	-	8,331
SANTA MARCELA	32	-	-	32	-	-	339	-	89	-	2,148	95	1,066	121	-	-	-	-	3,857
TOTAL	160	37	960	1,158	286	15	18,386	892	11,673	1,770	9,343	739	5,904	1,099	11	-	-	-	50,119

Note: Delivery of cacao planting materials must be started on the onset of rainy season.

AGRONOMIC REQUIREMENT OF CACAO PRODUCTION

LAND UTILIZATION TYPE	SUITABILITY RATING	SLOPE (%)	SOIL DEPTH (cm)	SOIL TEXTURE	SOIL DRAINAGE	SOIL REACTION (pH)	INHERENT FERTILITY	FLOODING CLASS	EROSION CLASS	ROCK OUTCROPS	ELEVATION (masl)	ANNUAL RAINFALL (mm)	CLIMATIC TYPE
	S1	<8	>100	CL, SiCL, SCL, SC, SiC, C, HC	WD,MWD	5.6 -7.2	high	none-slight	none-slight	none-few	<1000	2001-4500	I, III, IV
Cacao	S2	8 - 30	50 - 100	FSL, L, SiL	SPD,PD	5.1 - 5.5 7.3 - 7.8	medium	moderate	moderate	common	1000-1500	1000-2000	I, II
	S3	>30	<50	S, LS, CSL, SL	VPD,ED	<5.0 - > 7.9	low	severe	severe	many	>1500	<1000 >4500	

	S3	>30	<50	S, LS, CSL, SL	VPD,ED	<5.0 -	> 7.9 low	severe	sev	rere	many	>1500		000 500	
SLOPE (%	%)		SOIL DRA	AINAGE		SOIL REA	ACTION (pH)	1	SOIL TE	EXTURE	'				
0 - 3	- level to gently sloping	3	ED	- excessively drained		< 4.5	- extremely acid		Coarse			Fi	ine		
3 - 8	- gently sloping to und	ulating	WD	- well drained		4.5 - 5.0	- very strongly acid	d	S	- sand		SO	2	- sandy clay	
8 - 18	- undulating to rolling		MWD	- moderately well drained		5.1 - 5.5	- strongly acid		LS	- loam	y sand	Si	C	- silty clay	
18 - 30	- rolling to moderately	steep	SPD	- somewhat poorly draine	d	5.6 - 6.0	- medium acid		CSL	- coars	se sandy loam	С		- clay	
30 - 50	- steep		PD	 poorly drained 		6.1 - 6.5	 slightly acid 		SL	- sandy	/ loam	Н	C	- heavy clay	
> 50	- very steep		VPD	 very poorly drained 		6.6 - 7.2	- neutral		Mediur	n					
						7.3 - 7.8	 mildly alkaline 		FSL	- fine s	andy loam				
SOIL DE	РТН (ст)		SURFACI	E IMPEDIMENT		7.9 - 8.4	- moderately alkali	ne	L	- loam					
0 - 30	 very shallow 		ROCK OU	TCROPS		> 8.5	- strongly alkaline		SiL	- silt lo	am				
30 - 50	- shallow		< 10%	- none - few					CL	- clay l	oam				
50 - 100	- moderately deep		10 - 30%	- common					SiCL	- silty (clay loam				
> 100	- deep to very deep		> 30%	- many					SCL	- sandy	z clay loam				

LAND LIMITATIONS DESCRIPTION AND COMBINATIONS

ELEVATION	SOIL DRAINAGE	SOIL DEPTH	SOIL EROSION
El2 - 1000m - 1500m	D2 - Somewhat poorly drained to poorly drained	Sh2 - Moderately deep (50 - 100cm)	E2 - Moderate erosion
El3 -> 1500m	D3 - Very poorly drained or excessively drained	Sh3 - Very shallow to shallow (< 50cm)	E3 - Severe erosion
SLOPE/TOPOGRAPHY	SOIL TEXTURE	ROCK OUTCROPS	FLOODING
T2 - Undulating to moderately steep T3 - Steep to very steep	Tc - Coarse texture	Rc2 - Common Rc3 - Many	F2 - Moderate seasonal floodingF3 - Severe seasonal flooding

CODE	LAND LIMITATION	CODE	LAND LIMITATION	CODE	LAND LIMITATION	CODE	LAND LIMITATION	CODE	LAND LIMITATION
1	E2-Sh2-Rc2	11	El3-Sh2-Rc3	21	T2-El2-E3	31	T3-E3	41	T3-El2-E3-Sh3-Rc2
2	El2	12	F2-D2	22	T2-El2-E3-Rc2	32	T3-E3-Rc2	42	T3-El2-E3-Sh3-Rc3
3	El2-E2-Sh2-Rc3	13	Sh2	23	T2-El2-E3-Rc3	33	T3-E3-Rc3	43	T3-El3
4	El2-E3-Sh2-Rc3	14	Sh2-Rc2	24	T2-El2-E3-Sh2-Rc2	34	T3-E3-Sh3-Rc2	44	T3-El3-E3
5	El2-Rc2	15	T2	25	T2-El2-E3-Sh2-Rc3	35	T3-E3-Sh3-Rc3	45	T3-El3-E3-Sh2-Rc3
6	El2-Sh2-Rc2	16	T2-E3	26	T2-El3-E3	36	T3-El2	46	T3-El3-E3-Sh3-Rc3
7	El2-Sh2-Rc3	17	T2-E3-Rc2	27	T2-El3-E3-Rc3	37	T3-El2-E3		
8	El3	18	T2-E3-Sh2-Rc2	28	T2-El3-E3-Sh2-Rc3	38	T3-El2-E3-Rc2		
9	El3-E3-Sh2-Rc3	19	T2-E3-Sh2-Rc3	29	T2-F2-D2	39	T3-El2-E3-Rc3		
10	El3-Sh2-Rc2	20	T2-El2	30	T3	40	T3-El2-E3-Sh2-Rc3		

CODE	LAND USE
2	Rice paddy, non-irrigated
4	Corn
81	Coffee
82	Cacao
85	Mango
116	Coconut
126	Grassland
134	Shrubs, unmanaged

SUITABILITY CLASSES:

Highly Suitable (S1) Land having no significant limitation to sustained application of a given use, or only minor limitations that will not significantly reduce productivity or benefits and will not raise inputs above an acceptable level.

Marginally Suitable (S3) Land having limitations which in aggregate are severe for sustained application of a given use and will so reduce productivity or benefits, or increase required inputs, that this expenditure will be only marginally justified.

Moderately Suitable (S2) Land having limitation which in aggregate are moderately severe for sustained application of a given use; the limitation will reduce productivity or benefits and increase required inputs to the extent that the overall advantage to be gained from the use, although still attractive, will be appreciably

inferior to that expected on class S1 land.

Not Suitable / Not Relevant Land having limitations which may be surmountable in time but which cannot be corrected with existing knowledge at currently acceptable cost; the limitations are so severe as to preclude successful sustained use of the land in the given manner. Existing forest, shrubland greater than 18% slope, irrigated paddy rice and miscellaneous land types such as built up areas, roads, etc are considered as not relevant.

CLIMATE TYPE


TYPE I: Two pronouced season, dry from November to April and **TYPE II**: No dry season with a very pronounced maximum rain wet during the rest of the year. Maximum rain period is from June to September

period from December to February. There is not a single dry month. Maximum monthly rainfall occurs during the period from March to May.

TYPE III: No very pronounced maximum rain period, with a dry season lasting only from one to three months, either during the period from December to February or from March to May. This type resembles Type I since it has a short dry season.

TYPE IV: Rainfall is more or less evenly distributed throughout the year. This type resembles Type II since it has no dry

Mostly in the Northeastern part of the Apayao lies on Type III climate classification and the rest of the province is Type I.



Source: PAGASA 2018, Climatological Normals (Rainfall), Philippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA), accessed 27 July 2018, https://www1.pagasa.dost.gov.ph/index.php/climate/climatological-normals>.

^{*}establishment of shade trees prior to planting of cacao.

