



Department of Agriculture  
BUREAU OF SOILS AND WATER MANAGEMENT  
"NATIONAL SOIL SAMPLING AND TESTING FOR FERTILITY AND CROP SUITABILITY ASSESSMENT"  
**FERTILIZER-GUIDE FOR RICE**  
**PROVINCE OF BUKIDNON**

Municipality	Fertilizer-Guide Group		Additional 10-20 bags/ha Organic Fertilizer is recommended for all options																					
			INBRED RICE									HYBRID RICE												
	WET SEASON						DRY SEASON						WET SEASON						DRY SEASON					
	Number	Description	Nutrient Requirement			Recommended Fertilizer Rate (bags/hectare)	Nutrient Requirement			Recommended Fertilizer Rate (bags/hectare)	Nutrient Requirement			Recommended Fertilizer Rate (bags/hectare)	Nutrient Requirement			Recommended Fertilizer Rate (bags/hectare)						
			N	P <sub>2</sub> O	K <sub>2</sub> O		N	P <sub>2</sub> O	K <sub>2</sub> O		N	P <sub>2</sub> O	K <sub>2</sub> O		N	P <sub>2</sub> O	K <sub>2</sub> O							
KITAOTAO	122	M-H-MH-ML	30	20	40	Option 1: (1st Application): 2.25 (14-14-14-); 0.75 (0-18-0); (3rd Application): 0.75 (46-0-0); 1.00 (0-0-60); Option 2: (1st Application): 0.75 (46-0-0); 2.25 (0-18-0); (3rd Application): 0.75 (46-0-0); 1.50 (0-0-60);	40	20	40	Option 1: (1st Application): 3.00 (14-14-14-); (3rd Application): 1.00 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 1.00 (46-0-0); 2.25 (0-18-0); (3rd Application): 1.00 (46-0-0); 1.50 (0-0-60);	50	30	50	Option 1: (1st Application): 3.75 (14-14-14-); 0.75 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.00 (0-0-60); Option 2: (1st Application): 1.25 (46-0-0); 3.50 (0-18-0); (3rd Application): 1.00 (46-0-0); 1.75 (0-0-60);	60	30	50	Option 1: (1st Application): 4.50 (14-14-14-); (2nd Application): 1.50 (46-0-0); (3rd Application): 0.75 (0-0-60); Option 2: (1st Application): 1.50 (46-0-0); 3.50 (0-18-0); (2nd Application): 0.75 (46-0-0); (3rd Application): 0.75 (46-0-0); 1.75 (0-0-60);						
KITAOTAO	106	M-MH-MH-ML	50	20	40	Option 1: (1st Application): 3.00 (14-14-14-); (3rd Application): 1.00 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 2.00 (16-20-0); (2nd Application): 0.75 (46-0-0); (3rd Application): 0.75 (46-0-0); 1.50 (0-0-60); Option 3: (1st Application): 1.25 (46-0-0); 2.25 (0-18-0); (2nd Application): 0.50 (46-0-0); (3rd Application): 0.50 (46-0-0); 1.50 (0-0-60);	60	20	40	Option 1: (1st Application): 3.00 (14-14-14-); (2nd Application): 0.75 (46-0-0); (3rd Application): 0.75 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 2.00 (16-20-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.50 (0-0-60); Option 3: (1st Application): 1.50 (46-0-0); 2.25 (0-18-0); (2nd Application): 0.75 (46-0-0); (3rd Application): 0.75 (46-0-0); 1.50 (0-0-60);	70	30	50	Option 1: (1st Application): 4.50 (14-14-14-); (2nd Application): 1.00 (46-0-0); (3rd Application): 0.75 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 3.00 (16-20-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.75 (0-0-60); Option 3: (1st Application): 1.50 (46-0-0); 3.50 (0-18-0); (2nd Application): 0.75 (46-0-0); (3rd Application): 0.75 (46-0-0); 1.75 (0-0-60);	80	30	50	Option 1: (1st Application): 4.50 (14-14-14-); (2nd Application): 1.09 (46-0-0); (3rd Application): 0.87 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 3.00 (16-20-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.75 (0-0-60); Option 3: (1st Application): 1.75 (46-0-0); 3.50 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.75 (0-0-60);						

Application schedule:  
1st - after last harrowing  
2nd - 10-14 DAT  
3rd - 20-25 DAT     DAT - days after transplanting

Fertilizer materials:  
Complete (14-14-14)Urea (46-0-0)  
Ammophos (16-20-0)  
Solophos (0-18-0)  
Muriate of Potash (0-0-60)